

# A SUM OF SQUARES APPROXIMATION OF NONNEGATIVE POLYNOMIALS

JEAN B. LASSERRE

**ABSTRACT.** We show that every real nonnegative polynomial  $f$  can be approximated as closely as desired by a sequence of polynomials  $\{f_\epsilon\}$  that are sums of squares. Each  $f_\epsilon$  has a simple et explicit form in terms of  $f$  and  $\epsilon$ . A special representation is also obtained for convex polynomials, nonnegative on a convex semi-algebraic set.

## 1. INTRODUCTION

The study of relationships between *nonnegative* and *sums of squares* (s.o.s.) polynomials, initiated by Hilbert, is of real practical importance in view of numerous potential applications, notably in polynomial programming. Indeed, checking whether a given polynomial is nonnegative is a NP-hard problem whereas checking it is s.o.s. reduces to solving a (convex) Semidefinite Programming (SDP) problem for which efficient algorithms are now available. For instance, recent results in real algebraic geometry, most notably by Schmüdgen [15], Putinar [12], Jacobi and Prestel [4], have provided s.o.s. representations of polynomials, positive on a compact semialgebraic set; the interested reader is referred to Prestel and Delzell [11] and Scheiderer [14] for a nice account of such results. This in turn has permitted to develop efficient SDP-relaxations in polynomial optimization (see e.g. Lasserre [5, 6, 7], Parrilo [9, 10], Schweighofer [16], and the many references therein).

So, back to a comparison between nonnegative and s.o.s. polynomials, on the negative side, Blekherman [3] has shown that if the degree is *fixed*, then the cone of nonnegative polynomials is much *larger* than that of s.o.s. However, on the positive side, a denseness result [2] states that the cone of s.o.s. polynomials is *dense* in the space of polynomials that are nonnegative on  $[-1, 1]^n$  (for the norm  $\|f\|_1 = \sum_\alpha |f_\alpha|$  whenever  $f$  is written  $\sum_\alpha f_\alpha x^\alpha$  in the usual canonical basis); see e.g. Berg [2, Theorem 5, p. 117]).

**Contribution.** We show that *every nonnegative* polynomial  $f$  is almost a s.o.s., namely we show that  $f$  can be approximated by a sequence of s.o.s.

---

1991 *Mathematics Subject Classification.* 11E25 12D15 13P05, 12Y05, 90C22, 90C25.

*Key words and phrases.* Real algebraic geometry; nonnegative polynomials; sums of squares; semidefinite relaxations.

polynomials  $\{f_\epsilon\}_\epsilon$ , in the specific form

$$(1.1) \quad f_\epsilon = f + \epsilon \sum_{k=0}^{r_\epsilon} \sum_{j=1}^n \frac{x_j^{2k}}{k!},$$

for some  $r_\epsilon \in \mathbb{N}$ , so that  $\|f - f_\epsilon\|_1 \rightarrow 0$  as  $\epsilon \downarrow 0$ .

This result is in the spirit of the previous denseness result. However we here provide in (1.1) an *explicit* converging approximation with a very specific (and simple) form; namely it suffices to slightly perturbate  $f$  by adding a small coefficient  $\epsilon > 0$  to each square monomial  $x_i^{2k}$  for all  $i = 1, \dots, n$  and all  $k = 1, \dots, r$ , with  $r$  sufficiently large.

To prove this result we combine

- (generalized) **Carleman's** sufficient condition for a moment sequence  $\mathbf{y} = \{y_\alpha\}$  to have a *representing measure*  $\mu$  (i.e., such that  $y_\alpha = \int x^\alpha d\mu$  for all  $\alpha \in \mathbb{N}^n$ ), and
- a **duality** result from convex optimization.

As a consequence, we may thus define a procedure to approximate the global minimum of a polynomial  $f$ . It consists in solving a sequence of SDP-relaxations which are simpler and easier to solve than those defined in Lasserre [5]. Finally, we also consider the case where  $f$  is a *convex* polynomial, nonnegative on a convex semi-algebraic set  $\mathbb{K}$  defined by (concave polynomial) inequalities  $g_j \geq 0$ . We show that the approximation  $f_\epsilon$  of  $f$ , defined in (1.1), has a certificate of positivity on  $\mathbb{K}$  (or a representation) similar to Putinar's s.o.s. representation [12], but in which the s.o.s. polynomial coefficients of the  $g_j$ 's now become simple nonnegative *scalars*, the Lagrange multipliers of a related convex optimization problem.

## 2. NOTATION AND DEFINITIONS

For a real symmetric matrix  $A$ , the notation  $A \succeq 0$  (resp.  $A \succ 0$ ) stands for  $A$  positive semidefinite (resp. positive definite). The sup-norm  $\sup_j |x_j|$  of a vector  $x \in \mathbb{R}^n$ , is denoted by  $\|x\|_\infty$ . Let  $\mathbb{R}[x_1, \dots, x_n]$  be the ring of real polynomials, and let

$$(2.1) \quad v_r(x) := (1, x_1, x_2, \dots, x_n, x_1^2, x_1x_2, \dots, x_1x_n, x_2^2, x_2x_3, \dots, x_n^r)$$

be the canonical basis for the  $\mathbb{R}$ -vector space  $\mathcal{A}_r$  of real polynomials of degree at most  $r$ , and let  $s(r)$  be its dimension. Similarly,  $v_\infty(x)$  denotes the canonical basis of  $\mathbb{R}[x_1, \dots, x_n]$  as a  $\mathbb{R}$ -vector space, denoted  $\mathcal{A}$ . So a vector in  $\mathcal{A}$  has always *finitely* many zeros.

Therefore, a polynomial  $p \in \mathcal{A}_r$  is written

$$x \mapsto p(x) = \sum_{\alpha} p_{\alpha} x^{\alpha} = \langle \mathbf{p}, v_r(x) \rangle, \quad x \in \mathbb{R}^n,$$

(where  $x^{\alpha} = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}$ ) for some vector  $\mathbf{p} = \{p_{\alpha}\} \in \mathbb{R}^{s(r)}$ , the vector of coefficients of  $p$  in the basis (2.1).

Extending  $\mathbf{p}$  with zeros, we can also consider  $\mathbf{p}$  as a vector indexed in the basis  $v_\infty(x)$  (i.e.  $\mathbf{p} \in \mathcal{A}$ ). If we equip  $\mathcal{A}$  with the usual scalar product  $\langle \cdot, \cdot \rangle$  of vectors, then for every  $p \in \mathcal{A}$ ,

$$p(x) = \sum_{\alpha \in \mathbb{N}^n} p_\alpha x^\alpha = \langle \mathbf{p}, v_\infty(x) \rangle, \quad x \in \mathbb{R}^n.$$

Given a sequence  $\mathbf{y} = \{y_\alpha\}$  indexed in the basis  $v_\infty(x)$ , let  $L_{\mathbf{y}} : \mathcal{A} \rightarrow \mathbb{R}$  be the linear functional

$$p \mapsto L_{\mathbf{y}}(p) := \sum_{\alpha \in \mathbb{N}^n} p_\alpha y_\alpha = \langle \mathbf{p}, \mathbf{y} \rangle.$$

Given a sequence  $\mathbf{y} = \{y_\alpha\}$  indexed in the basis  $v_\infty(x)$ , the *moment* matrix  $M_r(\mathbf{y}) \in \mathbb{R}^{s(r) \times s(r)}$  with rows and columns indexed in the basis  $v_r(x)$  in (2.1), satisfies

$$[M_r(\mathbf{y})(1, j) = y_\alpha \text{ and } M_r(\mathbf{y})(i, 1) = y_\beta] \Rightarrow M_r(\mathbf{y})(i, j) = y_{\alpha+\beta}.$$

For instance, with  $n = 2$ ,

$$M_2(\mathbf{y}) = \begin{bmatrix} y_{00} & y_{10} & y_{01} & y_{20} & y_{11} & y_{02} \\ y_{10} & y_{20} & y_{11} & y_{30} & y_{21} & y_{12} \\ y_{01} & y_{11} & y_{02} & y_{21} & y_{12} & y_{03} \\ y_{20} & y_{30} & y_{21} & y_{40} & y_{31} & y_{22} \\ y_{11} & y_{21} & y_{12} & y_{31} & y_{22} & y_{13} \\ y_{02} & y_{12} & y_{03} & y_{22} & y_{13} & y_{04} \end{bmatrix}.$$

A sequence  $\mathbf{y} = \{y_\alpha\}$  has a *representing* measure  $\mu_{\mathbf{y}}$  if

$$(2.2) \quad y_\alpha = \int_{\mathbb{R}^n} x^\alpha d\mu_{\mathbf{y}}, \quad \forall \alpha \in \mathbb{N}^n.$$

In this case one also says that  $\mathbf{y}$  is a *moment sequence*. In addition, if  $\mu_{\mathbf{y}}$  is unique then  $\mathbf{y}$  is said to be a *determinate* moment sequence.

The matrix  $M_r(\mathbf{y})$  defines a bilinear form  $\langle \cdot, \cdot \rangle_{\mathbf{y}}$  on  $\mathcal{A}_r$ , by

$$\langle q, p \rangle_{\mathbf{y}} := \langle \mathbf{q}, M_r(\mathbf{y})\mathbf{p} \rangle = L_{\mathbf{y}}(qp), \quad q, p \in \mathcal{A}_r,$$

and if  $\mathbf{y}$  has a *representing* measure  $\mu_{\mathbf{y}}$  then

$$(2.3) \quad \langle \mathbf{q}, M_r(\mathbf{y})\mathbf{q} \rangle = \int_{\mathbb{R}^n} q(x)^2 d\mu_{\mathbf{y}}(x) \geq 0,$$

so that  $M_r(\mathbf{y}) \succeq 0$ .

Next, given a sequence  $\mathbf{y} = \{y_\alpha\}$  indexed in the basis  $v_\infty(x)$ , let  $y_{2k}^{(i)} := L_{\mathbf{y}}(x_i^{2k})$  for every  $i = 1, \dots, n$  and every  $k \in \mathbb{N}$ . That is,  $y_{2k}^{(i)}$  denotes the element in the sequence  $\mathbf{y}$ , corresponding to the monomial  $x_i^{2k}$ .

Of course not every sequence  $\mathbf{y} = \{y_\alpha\}$  has a representing measure  $\mu_{\mathbf{y}}$  as in (2.2). However, there exists a *sufficient* condition to ensure that it is the case. The following result stated in Berg [2, Theorem 5, p. 117] is from Nussbaum [8], and is re-stated here, with our notation.

**Theorem 2.1.** *Let  $\mathbf{y} = \{y_\alpha\}$  be an infinite sequence such that  $M_r(\mathbf{y}) \succeq 0$  for all  $r = 0, 1, \dots$ . If*

$$(2.4) \quad \sum_{k=0}^{\infty} (\mathbf{y}_{2k}^{(i)})^{-1/2k} = \infty, \quad i = 1, \dots, n,$$

*then  $\mathbf{y}$  is a determinate moment sequence.*

The condition (2.4) in Theorem 2.1 is called *Carleman's condition* as it extends to the multivariate case the original Carleman's sufficient condition given for the univariate case.

### 3. PRELIMINARIES

Let  $B_M$  be the closed ball

$$(3.1) \quad B_M = \{x \in \mathbb{R}^n \mid \|x\|_\infty \leq M\}.$$

**Proposition 3.1.** *Let  $f \in \mathbb{R}[x_1, \dots, x_n]$  be such that  $-\infty < f^* := \inf_x f(x)$ . Then, for every  $\epsilon > 0$  there is some  $M_\epsilon \in \mathbb{N}$  such that*

$$f_M^* := \inf_{x \in B_M} f(x) < f^* + \epsilon, \quad \forall M \geq M_\epsilon.$$

*Equivalently,  $f_M^* \downarrow f^*$  as  $M \rightarrow \infty$ .*

*Proof.* Suppose it is false. That is, there is some  $\epsilon_0 > 0$  and an infinite sequence  $\{M_k\} \subset \mathbb{N}$ , with  $M_k \rightarrow \infty$ , such that  $f_{M_k}^* \geq f^* + \epsilon_0$  for all  $k$ . But let  $x_0 \in \mathbb{R}^n$  be such that  $f(x_0) < f^* + \epsilon_0$ . With any  $M_k \geq \|x_0\|_\infty$ , one obtains the contradiction  $f^* + \epsilon_0 \leq f_{M_k}^* \leq f(x_0) < f^* + \epsilon_0$ .  $\square$

To prove our main result (Theorem 4.1 below), we first introduce the following related optimization problems.

$$(3.2) \quad \mathbb{P} : \quad f^* := \inf_{x \in \mathbb{R}^n} f(x),$$

and for  $0 < M \in \mathbb{N}$ ,

$$(3.3) \quad \mathcal{P}_M : \quad \inf_{\mu \in \mathcal{P}(\mathbb{R}^n)} \left\{ \int f d\mu \mid \int \sum_{i=1}^n e^{x_i^2} d\mu \leq n e^{M^2} \right\},$$

where  $\mathcal{P}(\mathbb{R}^n)$  is the space of probability measures on  $\mathbb{R}^n$ . The respective optimal values of  $\mathbb{P}$  and  $\mathcal{P}_M$  are denoted  $\inf \mathbb{P} = f^*$  and  $\inf \mathcal{P}_M$ , or  $\min \mathbb{P}$  and  $\min \mathcal{P}_M$  if the minimum is attained.

**Proposition 3.2.** *Let  $f \in \mathbb{R}[x_1, \dots, x_n]$  be such that  $-\infty < f^* := \inf_x f(x)$ , and consider the two optimization problems  $\mathbb{P}$  and  $\mathcal{P}_M$  defined in (3.2) and (3.3) respectively. Then,  $\inf \mathcal{P}_M \downarrow f^*$  as  $M \rightarrow \infty$ . If  $f$  has a global minimizer  $x^* \in \mathbb{R}^n$ , then  $\min \mathcal{P}_M = f^*$  whenever  $M \geq \|x^*\|_\infty$ .*

*Proof.* Let  $\mu \in \mathcal{P}(\mathbb{R}^n)$  be admissible for  $\mathcal{P}_M$ . As  $f \geq f^*$  on  $\mathbb{R}^n$  then it follows immediately that  $\int f d\mu \geq f^*$ , and so,  $\inf \mathcal{P}_M \geq f^*$  for all  $M$ .

As  $B_M$  is closed and bounded, it is compact and so, with  $f_M^*$  as in Proposition 3.1, there is some  $\hat{x} \in B_M$  such that  $f(\hat{x}) = f_M^*$ . In addition let  $\mu \in \mathcal{P}(\mathbb{R}^n)$  be the Dirac probability measure at the point  $\hat{x}$ . As  $\|\hat{x}\|_\infty \leq M$ ,

$$\int \sum_{i=1}^n e^{x_i^2} d\mu = \sum_{i=1}^n e^{(\hat{x}_i)^2} \leq ne^{M^2},$$

so that  $\mu$  is an admissible solution of  $\mathcal{P}_M$  with value  $\int f d\mu = f(\hat{x}) = f_M^*$ , which proves that  $\inf \mathcal{P}_M \leq f_M^*$ . This latter fact, combined with Proposition 3.1 and with  $f^* \leq \inf \mathcal{P}_M$ , implies  $\inf \mathcal{P}_M \downarrow f^*$  as  $M \rightarrow \infty$ , the desired result. The final statement is immediate by taking as feasible solution for  $\mathcal{P}_M$ , the Dirac probability measure at the point  $x^* \in B_M$  (with  $M \geq \|x^*\|_\infty$ ). As its value is now  $f^*$ , it is also optimal, and so,  $\mathcal{P}_M$  is solvable with optimal value  $\min \mathcal{P}_M = f^*$ .  $\square$

Proposition 3.2 provides a rationale for introducing the following Semi-definite Programming (SDP) problems. Let  $2r_f$  be the degree of  $f$  and for every  $r_f \leq r \in \mathbb{N}$ , consider the SDP problem

$$(3.4) \quad \mathbb{Q}_r \left\{ \begin{array}{ll} \min_{\mathbf{y}} L_{\mathbf{y}}(f) (= \sum_{\alpha} f_{\alpha} y_{\alpha}) & \\ \text{s.t. } M_r(\mathbf{y}) & \succeq 0 \\ \sum_{k=0}^r \sum_{i=1}^n \frac{y_{2k}^{(i)}}{k!} & \leq ne^{M^2}, \\ y_0 & = 1, \end{array} \right.$$

and its associated *dual* SDP problem

$$(3.5) \quad \mathbb{Q}_r^* \left\{ \begin{array}{ll} \max_{\lambda \geq 0, \gamma, q} \gamma - ne^{M^2} \lambda & \\ \text{s.t. } f - \gamma & = q - \lambda \sum_{k=0}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} \\ q & \text{s.o.s. of degree } \leq 2r, \end{array} \right.$$

with respective optimal values  $\inf \mathbb{Q}_r$  and  $\sup \mathbb{Q}_r^*$  (or  $\min \mathbb{Q}_r$  and  $\max \mathbb{Q}_r^*$  if the optimum is attained, in which case the problems are said to be solvable). For more details on SDP theory, the interested reader is referred to the survey paper [17].

The SDP problem  $\mathbb{Q}_r$  is a relaxation of  $\mathcal{P}_M$ , and we next show that in fact

- $\mathbb{Q}_r$  is solvable for all  $r \geq r_0$ ,
- its optimal value  $\min \mathbb{Q}_r \rightarrow \inf \mathcal{P}_M$  as  $r \rightarrow \infty$ , and
- $\mathbb{Q}_r^*$  is also solvable with same optimal value as  $\mathbb{Q}_r$ , for every  $r \geq r_f$ .

This latter fact will be crucial to prove our main result in the next section. Let  $l_\infty$  (resp.  $l_1$ ) be the Banach space of bounded (resp. summable) infinite sequences with the sup-norm (resp. the  $l_1$ -norm).

**Theorem 3.3.** *Let  $f \in \mathbb{R}[x_1, \dots, x_n]$  be of degree  $2r_f$ , with global minimum  $f^* > -\infty$ , and let  $M > 0$  be fixed. Then :*

- (i) *For every  $r \geq r_f$ ,  $\mathbb{Q}_r$  is solvable, and  $\min \mathbb{Q}_r \uparrow \inf \mathcal{P}_M$  as  $r \rightarrow \infty$ .*
- (ii) *Let  $\mathbf{y}^{(r)} = \{y_\alpha^{(r)}\}$  be an optimal solution of  $\mathbb{Q}_r$  and complete  $\mathbf{y}^{(r)}$  with zeros to make it an element of  $l_\infty$ . Every (pointwise) accumulation point  $\mathbf{y}^*$  of the sequence  $\{\mathbf{y}^{(r)}\}_{r \in \mathbb{N}}$  is a determinate moment sequence, that is,*

$$(3.6) \quad y_\alpha^* = \int_{\mathbb{R}^n} x^\alpha d\mu^*, \quad \alpha \in \mathbb{N}^n,$$

*for a unique probability measure  $\mu^*$ , and  $\mu^*$  is an optimal solution of  $\mathcal{P}_M$ .*

- (iii) *For every  $r \geq r_f$ ,  $\max \mathbb{Q}_r^* = \min \mathbb{Q}_r$ .*

For a proof see §5.1.

So, one can approximate the optimal value  $f^*$  of  $\mathbb{P}$  as closely as desired, by solving SDP-relaxations  $\{\mathbb{Q}_r\}$  for sufficiently large values of  $r$  and  $M$ . Indeed,  $f^* \leq \inf \mathcal{P}_M \leq f_M^*$ , with  $f_M^*$  as in Proposition 3.1. Therefore, let  $\epsilon > 0$  be fixed, arbitrary. By Proposition 3.2, we have  $f^* \leq \inf \mathcal{P}_M \leq f^* + \epsilon$  provided that  $M$  is sufficiently large. Next, by Theorem 3.3(i), one has  $\inf \mathbb{Q}_r \geq \inf \mathcal{P}_M - \epsilon$  provided that  $r$  is sufficiently large, in which case, we finally have  $f^* - \epsilon \leq \inf \mathbb{Q}_r \leq f^* + \epsilon$ .

Notice that the SDP-relaxation  $\mathbb{Q}_r$  in (3.4) is simpler than the one defined in Lasserre [5]. Both have the same variables  $\mathbf{y} \in \mathbb{R}^{s(r)}$ , but the former has *one* SDP constraint  $M_r(\mathbf{y}) \succeq 0$  and one scalar inequality (as one substitutes  $y_0$  with 1) whereas the latter has the same SDP constraint  $M_r(\mathbf{y}) \succeq 0$  and one additional SDP constraint  $M_{r-1}(\theta\mathbf{y}) \succeq 0$  for the localizing matrix associated with the polynomial  $x \mapsto \theta(x) = M^2 - \|x\|^2$ . This results in a significant simplification.

#### 4. SUM OF SQUARES APPROXIMATION

Let  $\mathcal{A}$  be equipped with the norm

$$f \mapsto \|f\|_1 := \sum_{\alpha \in \mathbb{N}^n} |f_\alpha|, \quad f \in \mathcal{A}.$$

**Theorem 4.1.** *Let  $f \in \mathbb{R}[x_1, \dots, x_n]$  be nonnegative with global minimum  $f^*$ , that is,*

$$0 \leq f^* \leq f(x), \quad x \in \mathbb{R}^n.$$

- (i) *There is some  $r_0 \in \mathbb{N}$ ,  $\lambda_0 \geq 0$  such that, for all  $r \geq r_0$  and  $\lambda \geq \lambda_0$ ,*

$$(4.1) \quad f + \lambda \sum_{k=0}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} \quad \text{is a sum of squares.}$$

(ii) For every  $\epsilon > 0$ , there is some  $r_\epsilon \in \mathbb{N}$  such that,

$$(4.2) \quad f_\epsilon := f + \epsilon \sum_{k=0}^{r_\epsilon} \sum_{j=1}^n \frac{x_j^{2k}}{k!} \quad \text{is a sum of squares.}$$

Hence,  $\|f - f_\epsilon\|_1 \rightarrow 0$  as  $\epsilon \downarrow 0$ .

For a proof see §5.2.

*Remark 4.2.* Theorem 4.1(ii) is a *denseness* result in the spirit of Theorem 5 in Berg [2, p. 117] which states that the cone of s.o.s. polynomials is dense (also for the norm  $\|f\|_1$ ) in the cone of polynomials that are nonnegative on  $[-1, 1]^n$ . However, notice that Theorem 4.1(ii) provides an *explicit* converging sequence  $\{f_\epsilon\}$  with a simple and very specific form.

We next consider the case of a convex polynomial, nonnegative on a convex semi-algebraic set. Given  $\{g_j\}_{j=1}^m \subset \mathbb{R}[x_1, \dots, x_n]$ , let  $\mathbb{K} \subset \mathbb{R}^n$  be the semi-algebraic set

$$(4.3) \quad \mathbb{K} := \{x \in \mathbb{R}^n \mid g_j(x) \geq 0, \quad j = 1, \dots, m\}.$$

**Corollary 4.3.** *Let  $\mathbb{K}$  be as in (4.3), where all the  $g_j$ 's are concave, and assume that Slater's condition holds, i.e., there exists  $x_0 \in \mathbb{K}$  such that  $g_j(x_0) > 0$  for all  $j = 1, \dots, m$ .*

*Let  $f \in \mathbb{R}[x_1, \dots, x_n]$  be convex, nonnegative on  $\mathbb{K}$ , and with a minimizer on  $\mathbb{K}$ , that is,  $f(x^*) \leq f(x)$  for all  $x \in \mathbb{K}$ , for some  $x^* \in \mathbb{K}$ . Then there exists a nonnegative vector  $\lambda \in \mathbb{R}^m$  such that for every  $\epsilon > 0$ , there is some  $r_\epsilon \in \mathbb{N}$  for which*

$$(4.4) \quad f + \epsilon \sum_{k=0}^{r_\epsilon} \sum_{i=1}^n \frac{x_i^{2k}}{k!} = f_0 + \sum_{j=1}^m \lambda_j g_j,$$

with  $f_0 \in \mathbb{R}[x_1, \dots, x_n]$  being a sum of squares. (Therefore, the degree of  $f_0$  is less than  $\max[2r_\epsilon, \deg f, \deg g_1, \dots, \deg g_m]$ .)

*Proof.* Consider the convex optimization problem  $f^* := \min\{f(x) \mid x \in \mathbb{K}\}$ . As  $f$  is convex,  $\mathbb{K}$  is a convex set and Slater's condition holds, the Karush-Kuhn-Tucker optimality condition holds. That is, there exists a *nonnegative* vector  $\lambda \in \mathbb{R}^m$  of Lagrange-KKT multipliers, such that

$$\nabla f(x^*) = \sum_{j=1}^m \lambda_j \nabla g_j(x^*); \quad \lambda_j g_j(x^*) = 0, \quad j = 1, \dots, m.$$

(See e.g. Rockafellar [13].) In other words,  $x^*$  is also a (global) minimizer of the convex Lagrangian  $L := f - \sum_{j=1}^m \lambda_j g_j$ . Then  $f^* = f(x^*) = L(x^*)$  is the (global) minimum of  $f$  on  $\mathbb{K}$ , as well as the global minimum of  $L$  on  $\mathbb{R}^n$ , i.e.,

$$(4.5) \quad f - \sum_{j=1}^m \lambda_j g_j - f^* \geq 0, \quad x \in \mathbb{R}^n.$$

As  $f \geq 0$  on  $\mathbb{K}$ ,  $f^* \geq 0$ , and so  $L \geq 0$  on  $\mathbb{R}^n$ . Then (4.4) follows from Theorem 4.1(ii), applied to the polynomial  $L$ .  $\square$

When  $\mathbb{K}$  is *compact* (and so,  $f$  has necessarily a minimizer  $x^* \in \mathbb{K}$ ), one may compare Corollary 4.3 with Putinar's representation [12] of polynomials, positive on  $\mathbb{K}$ . When  $f$  is nonnegative on  $\mathbb{K}$  (compact), and with

$$(4.6) \quad f_\epsilon := f + \epsilon \sum_{k=0}^{r_\epsilon} \sum_{i=1}^n \frac{x_i^{2k}}{k!},$$

one may rewrite (4.4) as

$$(4.7) \quad f_\epsilon = f_0 + \sum_{j=1}^m \lambda_j g_j,$$

which is indeed a *certificate* of positivity of  $f_\epsilon$  on  $\mathbb{K}$ . In fact, as  $f_\epsilon > 0$  on  $\mathbb{K}$ , (4.7) can be seen as a special form of Putinar's s.o.s. representation, namely

$$(4.8) \quad f_\epsilon = q_0 + \sum_{j=1}^m q_j g_j, \quad \text{with } q_0, \dots, q_m \text{ s.o.s.}$$

(which holds under an additional assumption on the  $g_j$ 's). So, in the convex compact case, and under Slater's condition, Corollary 4.3 states that if  $f \geq 0$  on  $\mathbb{K}$ , then its approximation  $f_\epsilon$  in (4.6), has the simplified Putinar representation (4.7), in which the s.o.s. coefficients  $\{q_j\}$  of the  $g_j$ 's in (4.8), become now simple nonnegative *scalars* in (4.7), namely, the Lagrange-KKT multipliers  $\{\lambda_j\}$ .

## 5. PROOFS

**5.1. Proof of Theorem 3.3.** We will prove (i) and (ii) together. We first prove that  $\mathbb{Q}_r$  is solvable. This is because the feasible set (which is closed) is compact. Indeed, the constraint

$$\sum_{k=0}^r \sum_{i=1}^n y_{2k}^{(i)} / k! \leq ne^{M^2}$$

implies that every diagonal element  $y_{2k}^{(i)}$  of  $M_r(\mathbf{y})$  is bounded by  $\tau_r := nr!e^{M^2}$ . By Lemma 6.2, this in turn implies that its diagonal elements (i.e.,  $y_{2\alpha}$ , with  $|\alpha| \leq r$ ) are *all* bounded by  $\tau_r$ .

This latter fact and again  $M_r(\mathbf{y}) \succeq 0$ , also imply that in fact *every* element of  $M_r(\mathbf{y})$  is bounded by  $\tau_r$ , that is,  $|y_\alpha| \leq \tau_r$  for all  $|\alpha| \leq 2r$ . Indeed, for a symmetric matrix  $A \succeq 0$ , every non diagonal element  $A_{ij}$  satisfies  $A_{ij}^2 \leq A_{ii}A_{jj}$  so that  $|A_{ij}| \leq \max_i A_{ii}$ .

Therefore the set of feasible solutions of  $\mathbb{Q}_r$  is a closed bounded subset of  $\mathbb{R}^{s(r)}$ , hence compact. As  $L_{\mathbf{y}}(f)$  is linear in  $\mathbf{y}$ , the infimum is attained at some feasible point. Thus, for all  $r \geq r_f$ ,  $\mathbb{Q}_r$  is solvable with optimal value



$\min \mathbb{Q}_r \leq \inf \mathcal{P}_M$ . The latter inequality is because the moment sequence  $\mathbf{y}$  associated with an arbitrary feasible solution  $\mu$  of  $\mathcal{P}_M$ , is obviously feasible for  $\mathbb{Q}_r$ , and with value  $L_{\mathbf{y}}(f) = \int f d\mu$ .

Next, as the sequence  $\{\min \mathbb{Q}_r\}_r$  is obviously monotone non decreasing, one has  $\min \mathbb{Q}_r \uparrow \rho^* \leq \inf \mathcal{P}_M$ , as  $r \rightarrow \infty$ . We have seen that every entry of  $M_r(\mathbf{y})$  is bounded by  $\tau_r$ , and this bound holds for all  $r \geq r_f$ . Moreover,  $M_r(\mathbf{y})$  is also a (north-west corner) submatrix of  $M_s(\mathbf{y})$  for every  $s > r$ . Indeed, whenever  $s > r$ , one may write

$$M_s(\mathbf{y}) = \left[ \begin{array}{c|c} M_r(\mathbf{y}) & B \\ \hline B' & C \end{array} \right]$$

for some appropriate matrices  $B$  and  $C$ . Therefore, for the same reasons, any feasible solution  $\mathbf{y}$  of  $\mathbb{Q}_s(\mathbf{y})$  satisfies  $|y_\alpha| \leq \tau_r$ , for all  $\alpha \in \mathbb{N}^n$  such that  $|\alpha| \leq 2r$ . Therefore, for every  $s \in \mathbb{N}$ , and every feasible solution  $\mathbf{y}$  of  $\mathbb{Q}_s$ , we have

$$|y_\alpha| \leq \tau_r, \quad \forall \alpha \in \mathbb{N}^n, \quad 2r - 1 \leq |\alpha| \leq 2r, \quad r = 1, \dots, s.$$

Thus, given  $\mathbf{y} = \{y_\alpha\}$ , denote by  $\hat{\mathbf{y}} = \{\hat{\mathbf{y}}_\alpha\}$  the new sequence obtained from  $\mathbf{y}$  by the scaling

$$\hat{\mathbf{y}}_\alpha := \mathbf{y}_\alpha / \tau_r \quad \forall \alpha \in \mathbb{N}^n, \quad 2r - 1 \leq |\alpha| \leq 2r, \quad r = 1, 2, \dots$$

So let  $\mathbf{y}^{(r)} = \{y_\alpha^{(r)}\}$  be an optimal solution of  $\mathbb{Q}_r$  and complete  $\mathbf{y}^{(r)}$  with zeros to make it an element of  $l_\infty$ . Hence, all the elements  $\hat{\mathbf{y}}^{(r)}$  are in the unit ball  $B_1$  of  $l_\infty$ , defined by

$$B_1 = \{\mathbf{y} = \{y_\alpha\} \in l_\infty \mid \|\mathbf{y}\|_\infty \leq 1\}.$$

By the Banach-Alaoglu Theorem, this ball is sequentially compact in the  $\sigma(l_\infty, l_1)$  (weak\*) topology of  $l_\infty$  (see e.g. Ash [1]). In other words, there exists an element  $\hat{\mathbf{y}}^* \in B_1$  and a subsequence  $\{r_k\} \subset \mathbb{N}$ , such that  $\hat{\mathbf{y}}^{(r_k)} \rightarrow \hat{\mathbf{y}}^*$  for the weak\* topology of  $l_\infty$ , that is, for all  $u \in l_1$ ,

$$(5.1) \quad \langle \hat{\mathbf{y}}^{(r_k)}, u \rangle \rightarrow \langle \hat{\mathbf{y}}^*, u \rangle, \quad \text{as } k \rightarrow \infty.$$

In particular, *pointwise* convergence holds, that is, for all  $\alpha \in \mathbb{N}^n$ ,

$$\hat{y}_\alpha^{(r_k)} \rightarrow \hat{y}_\alpha^*, \quad \text{as } k \rightarrow \infty,$$

and so, defining  $\mathbf{y}^*$  from  $\hat{\mathbf{y}}^*$  by

$$\mathbf{y}_\alpha^* = \tau_r \hat{\mathbf{y}}_\alpha^*, \quad \forall \alpha \in \mathbb{N}^n, \quad 2r - 1 \leq |\alpha| \leq 2r, \quad r = 1, 2, \dots$$

one also obtains the pointwise convergence

$$(5.2) \quad \text{for all } \alpha \in \mathbb{N}^n, \quad y_\alpha^{(r_k)} \rightarrow y_\alpha^*, \quad \text{as } k \rightarrow \infty.$$

We next prove that  $\mathbf{y}^*$  is the moment sequence of an optimal solution  $\mu^*$  of problem  $\mathcal{P}_M$ . From the pointwise convergence (5.2), we immediately get

$M_r(\mathbf{y}^*) \succeq 0$  for all  $r \geq r_f$ , because  $M_r(\mathbf{y})$  belongs to the cone of positive semidefinite matrices of size  $s(r)$ , which is closed. Next, and again by pointwise convergence, for every  $s \in \mathbb{N}$ ,

$$\sum_{j=0}^s \sum_{i=1}^n (y^*)_{2j}^{(i)} / j! = \lim_{k \rightarrow \infty} \sum_{j=0}^s \sum_{i=1}^n (y^{(r_k)})_{2j}^{(i)} / j! \leq ne^{M^2},$$

and so, by the Monotone Convergence Theorem

$$(5.3) \quad \sum_{j=0}^{\infty} \sum_{i=1}^n (y^*)_{2j}^{(i)} / j! = \lim_{s \rightarrow \infty} \sum_{j=0}^s \sum_{i=1}^n (y^*)_{2j}^{(i)} / j! \leq ne^{M^2}.$$

But (5.3) implies that  $\mathbf{y}^*$  satisfies Carleman's condition (2.4) is satisfied. Indeed, from (5.3), for all  $i = 1, \dots, n$ , we have  $(y^*)_{2k}^{(i)} < \rho k!$  for all  $k \in \mathbb{N}$ , and so, as  $k! \leq k^k = \sqrt{k}^{2k}$ ,

$$[(y^*)_{2k}^{(i)}]^{-1/2k} > (\rho)^{-1/2k} / \sqrt{k},$$

which in turn implies

$$\sum_{k=0}^{\infty} [(y^*)_{2k}^{(i)}]^{-1/2k} > \sum_{k=0}^{\infty} \frac{\rho^{-1/2k}}{\sqrt{k}} = +\infty.$$

Hence, by Theorem 2.1,  $\mathbf{y}^*$  is a determinate moment sequence, that is, there exists a unique measure  $\mu^*$  on  $\mathbb{R}^n$ , such that

$$y_{\alpha}^* = \int_{\mathbb{R}^n} x^{\alpha} d\mu^*, \quad \alpha \in \mathbb{N}^n.$$

By (5.3),

$$\int \sum_{i=1}^n e^{x_i^2} d\mu^* = \sum_{j=0}^{\infty} \sum_{i=1}^n (y^*)_{2j}^{(i)} / j! \leq ne^{M^2},$$

which proves that  $\mu^*$  is admissible for  $\mathcal{P}_M$ .

But then, again by the pointwise convergence (5.2) of  $\mathbf{y}^{(r_k)}$  to  $\mathbf{y}^*$ , we get  $L_{\mathbf{y}^{(r_k)}}(f) \rightarrow L_{\mathbf{y}^*}(f) = \int f d\mu^*$  as  $k \rightarrow \infty$ , which, in view of  $L_{\mathbf{y}^{(r_k)}}(f) \leq \inf \mathcal{P}_M$  for all  $k$ , implies

$$\int f d\mu^* = L_{\mathbf{y}^*}(f) \leq \inf \mathcal{P}_M.$$

But this proves that  $\mu^*$  is an optimal solution of  $\mathcal{P}_M$  because  $\mu^*$  is admissible for  $\mathcal{P}_M$  with value  $\int f d\mu^* \leq \inf \mathcal{P}_M$ . As the converging subsequence  $\{r_k\}$  was arbitrary, it is true for every limit point. Hence, we have proved (i) and (ii).

(iii) Let  $\mathbf{y}$  be the moment sequence associated with the probability measure  $\mu$  on the ball

$$B_{M/2} = \{\mathbf{y} = \{y_{\alpha}\} \in l_{\infty} \mid \|\mathbf{y}\|_{\infty} \leq M/2\},$$

with uniform distribution. That is,

$$\mu(B) = M^{-n} \int_{B \cap B_{M/2}} dx, \quad B \in \mathcal{B},$$

where  $\mathcal{B}$  is the sigma-algebra of Borel subsets of  $\mathbb{R}^n$ .

As  $\mu$  has a continuous density  $f_\mu > 0$  on  $B_{M/2}$ , it follows easily that  $M_r(\mathbf{y}) \succ 0$  for all  $r \geq r_f$ . In addition,

$$\sum_{k=0}^r \sum_{i=1}^n y_{2k}^{(i)} / k! < \int \sum_{i=1}^n e^{x_i^2} d\mu < ne^{M^2},$$

so that  $\mathbf{y}$  is a strictly admissible solution for  $\mathbb{Q}_r$ . Hence, the SDP problem  $\mathbb{Q}_r$  satisfies Slater's condition, and so, there is no duality gap between  $\mathbb{Q}_r$  and  $\mathbb{Q}_r^*$ , and  $\mathbb{Q}_r^*$  is solvable if  $\inf \mathbb{Q}_r$  is finite; see e.g. Vandenberghe and Boyd [17]. Thus,  $\mathbb{Q}_r^*$  is solvable because we proved that  $\mathbb{Q}_r$  is solvable. In other words,  $\sup \mathbb{Q}_r^* = \max \mathbb{Q}_r^* = \min \mathbb{Q}_r$ , the desired result.  $\square$

**5.2. Proof of Theorem 4.1.** It suffices to prove (i) and (ii) for the case  $f^* > 0$ . Indeed, if  $f^* = 0$  take  $\epsilon > 0$  arbitrary, fixed. Then  $f + n\epsilon \geq f_\epsilon^* = f^* + n\epsilon > 0$  and so, suppose that (4.1) holds for  $f + n\epsilon$  (for some  $r_0, \lambda_0$ ). In particular, pick  $\lambda \geq \lambda_0 + \epsilon$ , so that

$$f + n\epsilon + (\lambda - \epsilon) \sum_{k=0}^{r_\lambda} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = q_\lambda,$$

(with  $q_\lambda$  s.o.s.), for  $r_\lambda \geq r_0$ . Equivalently,

$$f + \lambda \sum_{k=0}^{r_\lambda} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = q_\lambda + \epsilon \sum_{k=1}^{r_\lambda} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = \hat{q}_\lambda,$$

where  $\hat{q}_\lambda$  is a s.o.s. Hence (4.1) also holds for  $f$  (with  $\lambda_0 + \epsilon$  in lieu of  $\lambda_0$ ).

Similarly, for (4.2). As  $f^* = 0$ ,  $f + n\epsilon > 0$  and so, suppose that (4.2) holds for  $f + n\epsilon$ . In particular,

$$f + n\epsilon + \epsilon \sum_{k=0}^{r_\epsilon} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = q_\epsilon,$$

(with  $q_\epsilon$  s.o.s.), for some  $r_\epsilon$ . Equivalently,

$$f + 2\epsilon \sum_{k=0}^{r_\epsilon} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = q_\epsilon + \epsilon \sum_{k=1}^{r_\epsilon} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = \hat{q}_\epsilon,$$

where  $\hat{q}_\epsilon$  is a s.o.s. Hence (4.2) also holds for  $f$ . Therefore, we will assume that  $f^* > 0$ .

(i) As  $f^* > 0$ , let  $M_0$  be such that  $f^* > 1/M_0$ , and fix  $M > M_0$ . Consider the SDP problem  $\mathbb{Q}_r^*$  defined in (3.5), associated with  $M$ . By Proposition 3.2,  $f^* \leq \inf \mathcal{P}_M$ . By Theorem 3.3,  $\max \mathbb{Q}_r^* = \min \mathbb{Q}_r \uparrow \inf \mathcal{P}_M \geq f^*$ . Therefore, there exists some  $r_M \geq r_f$  such that  $\max \mathbb{Q}_{r_M}^* \geq f^* - 1/M > 0$ .

That is, if  $(q_M, \lambda_M, \gamma_M)$  is an optimal solution of  $\mathbb{Q}_{r_M}^*$ , then  $\gamma_M - n\lambda_M e^{M^2} \geq f^* - 1/M > 0$ . In addition,

$$f - \gamma_M = q_M - \lambda_M \sum_{k=0}^{r_M} \sum_{j=1}^n \frac{x_j^{2k}}{k!},$$

that we rewrite

$$(5.4) \quad f - (\gamma_M - n\lambda_M e^{M^2}) = q_M + \lambda_M \left( n e^{M^2} - \sum_{k=0}^{r_M} \sum_{j=1}^n \frac{x_j^{2k}}{k!} \right).$$

Equivalently,

$$f + \lambda_M \sum_{k=0}^{r_M} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = q_M + n\lambda_M e^{M^2} + (\gamma_M - n\lambda_M e^{M^2}).$$

Define  $\hat{q}_M$  to be the s.o.s. polynomial

$$\hat{q}_M := q_M + n\lambda_M e^{M^2} + (\gamma_M - n\lambda_M e^{M^2}),$$

so that we obtain

$$(5.5) \quad f + \lambda_M \sum_{k=0}^{r_M} \sum_{j=1}^n \frac{x_j^{2k}}{k!} = \hat{q}_M,$$

the desired result.

If we now take  $r > r_M$  and  $\lambda \geq \lambda_M$  we also have

$$\begin{aligned} f + \lambda \sum_{k=0}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} &= f + \lambda_M \sum_{k=0}^{r_M} \sum_{j=1}^n \frac{x_j^{2k}}{k!} \\ &\quad + \lambda_M \sum_{k=r_M+1}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} + (\lambda - \lambda_M) \sum_{k=0}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} \\ &= \hat{q}_M + \lambda_M \sum_{k=r_M+1}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} + (\lambda - \lambda_M) \sum_{k=0}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} \\ &= \hat{\hat{q}}_M, \end{aligned}$$

that is,

$$(5.6) \quad f + \lambda \sum_{k=0}^r \sum_{j=1}^n \frac{x_j^{2k}}{k!} = \hat{\hat{q}}_M,$$

where  $\hat{\hat{q}}_M$  is a s.o.s. polynomial, the desired result.  $\square$

(ii) Let  $M$  be as in (i) above. Evaluating (5.4) at  $x = 0$ , and writing  $f(0) = f^* - f^* + f^*$ , yields

$$f(0) - f^* + f^* - (\gamma_M - n\lambda_M e^{M^2}) = q_M(0) + n\lambda_M (e^{M^2} - 1),$$

and as  $1/M > f^* - (\gamma_M - n\lambda_M e^{M^2})$ ,

$$\lambda_M \leq \frac{1/M + f(0) - f^*}{n(e^{M^2} - 1)}.$$

Now, letting  $M \rightarrow \infty$ , yields  $\lambda_M \rightarrow 0$ .

Now, let  $\epsilon > 0$  be fixed, arbitrary. There is some  $M > M_0$  such that  $\lambda_M \leq \epsilon$  in (5.5). Therefore, (4.2) is just (5.6) with  $\lambda := \epsilon > \lambda_M$  and  $r = r_\epsilon \geq r_M$ . Finally, from this, we immediately have

$$\|f - f_\epsilon\|_1 \leq \epsilon \sum_{i=1}^n \sum_{k=0}^{\infty} \frac{1}{k!} = \epsilon n e \rightarrow 0, \quad \text{as } \epsilon \downarrow 0.$$

□

## 6. APPENDIX

In this section we derive two auxiliary results that are helpful in the proofs of Theorem 3.3 and Theorem 4.1 in §5.

**Lemma 6.1.** *Let  $n = 2$  and let  $\mathbf{y}$  be a sequence indexed in the basis (2.1), and such that  $M_r(\mathbf{y}) \succeq 0$ . Then all the diagonal entries of  $M_r(\mathbf{y})$  are bounded by  $\tau_r := \max_{k=1, \dots, r} \max[y_{2k,0}, y_{0,2k}]$ .*

*Proof.* It suffices to prove that all the entries  $y_{2\alpha, 2\beta}$  with  $|\alpha + \beta| = r$  are bounded by  $s_r := \max[y_{2r,0}, y_{0,2r}]$ , and repeat the argument for entries  $y_{2\alpha, 2\beta}$  with  $|\alpha + \beta| = r-1, r-2$ , etc ... Then, take  $\tau_r := \max_{k=1, \dots, r} s_k$ . So, consider the odd case  $r = 2p+1$ , and the even case  $r = 2p$ .

- The odd case  $r = 2p+1$ . Let  $\Gamma := \{(2\alpha, 2\beta) \mid \alpha + \beta = r, \alpha \neq 0\}$ , and notice that

$$\Gamma = \{(2r-2k, 2k) \mid k = 1, \dots, r-1\} = \Gamma_1 \cup \Gamma_2$$

with

$$\Gamma_1 := \{(r, 0) + (r-2k, 2k), \mid k = 1, \dots, p\},$$

and

$$\Gamma_2 := \{(0, r) + (2j, r-2j), \mid j = 1, \dots, p\}.$$

Therefore, consider the two rows (and columns) corresponding to the indices  $(r, 0)$  and  $(r-2k, 2k)$ , or  $(0, r)$  and  $(2j, r-2j)$ . In view of  $M_r(\mathbf{y}) \succeq 0$ , one has

$$(6.1) \quad \begin{cases} y_{2r,0} \times y_{2r-4k,4k} & \geq (y_{2r-2k,2k})^2, & k = 1, \dots, p, \\ y_{0,2r} \times y_{4j,2r-4j} & \geq (y_{2j,2r-2j})^2, & j = 1, \dots, p. \end{cases}$$

Thus, let  $s := \max\{y_{2\alpha, 2\beta} \mid \alpha + \beta = r, \alpha \neq 0\}$ , so that either  $s = y_{2r-2k^*, 2k^*}$  for some  $1 \leq k^* \leq p$ , or  $s = y_{2j^*, 2r-2j^*}$  for some  $1 \leq j^* \leq p$ . But then, in view of (6.1), and with  $s_r := \max[y_{2r,0}, y_{0,2r}]$ ,

$$s_r \times s \geq y_{2r,0} \times y_{2r-4k^*, 4k^*} \geq (y_{2r-2k^*, 2k^*})^2 = s^2,$$

or,

$$s_r \times s \geq y_{0,2r} \times y_{4j^*, 2r-4j^*} \geq (y_{2j^*, 2r-2j^*})^2 = s^2,$$

so that  $s \leq s_r$ , the desired result.

- The even case  $r = 2p$ . Again, the set  $\Gamma := \{(2\alpha, 2\beta) \mid \alpha + \beta = r, \alpha \neq 0\}$  can be written  $\Gamma = \Gamma_1 \cup \Gamma_2$ , with

$$\Gamma_1 := \{(r, 0) + (r - 2k, 2k), \mid k = 1, \dots, p\},$$

and

$$\Gamma_2 := \{(0, r) + (2j, r - 2j), \mid j = 1, \dots, p\}.$$

The only difference with the odd case is that  $\Gamma_1 \cap \Gamma_2 = (p, p) \neq \emptyset$ . But the rest of the proof is the same as in the odd case.  $\square$

**Lemma 6.2.** *Let  $r \in \mathbb{N}$  be fixed, and let  $\mathbf{y}$  be a sequence such that the associated moment matrix  $M_r(\mathbf{y})$  is positive semidefinite, i.e.,  $M_r(\mathbf{y}) \succeq 0$ . Assume that there is some  $\tau_r \in \mathbb{R}$  such that the diagonal elements  $\{y_{2k}^{(i)}\}$  satisfy  $y_{2k}^{(i)} \leq \tau_r$ , for all  $k = 1, \dots, r$ , and all  $i = 1, \dots, n$ .*

*Then, the diagonal elements of  $M_r(\mathbf{y})$  are all bounded by  $\tau_r$  (i.e.,  $y_{2\alpha} \leq \tau_r$  for all  $\alpha \in \mathbb{N}^n$ , with  $|\alpha| \leq r$ ).*

*Proof.* The proof is by induction on the the number  $n$  of variables. By our assumption it is true for  $n = 1$ , and by Lemma 6.1, it is true for  $n = 2$ . Thus, suppose it is true for  $k = 1, 2, \dots, n - 1$  variables and consider the case of  $n$  variables (with  $n > 3$ ).

By our induction hypothesis, it is true for all elements  $y_{2\alpha}$  where at least one index, say  $\alpha_i$ , is zero ( $\alpha_i = 0$ ). Indeed, the submatrix  $A_r^{(i)}(\mathbf{y})$  of  $A_r(\mathbf{y})$ , obtained from  $A_r(\mathbf{y})$  by deleting all rows and columns corresponding to indices  $\alpha \in \mathbb{N}^n$  in the basis (2.1), with  $\alpha_i > 0$ , is a moment matrix of order  $r$ , with  $n - 1$  variables  $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n$ . Hence, by a permutation of rows and columns, we can write

$$A_r(\mathbf{y}) = \left[ \begin{array}{c|c} A_r^{(i)}(\mathbf{y}) & B \\ \hline B' & C \end{array} \right],$$

for some appropriate matrices  $B$  and  $C$ . In particular, all elements  $y_{2\alpha}$  with  $\alpha_i = 0$ , are diagonal elements of  $A_r^{(i)}(\mathbf{y})$ . In addition, its diagonal elements  $y_{2k}^{(j)}$ ,  $j \neq i$ , are all bounded by  $\tau_r$ . And of course,  $A_r^{(i)}(\mathbf{y}) \succeq 0$ . Therefore, by our induction hypothesis, all its diagonal elements are bounded by  $\tau_r$ . As  $i$  was arbitrary, we conclude that all elements  $y_{2\alpha}$  with at least one index being zero, are all bounded by  $\tau_r$ .

We next prove it is true for an arbitrary element  $y_{2\alpha}$  with  $|\alpha| \leq r$  and  $\alpha > 0$ , i.e.,  $\alpha_j \geq 1$  for all  $j = 1, \dots, n$ . With no loss of generality, we assume that  $\alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_n$ .

Consider the two elements  $y_{2\alpha_1, 0, \beta}$  and  $y_{0, 2\alpha_2, \gamma}$ , with  $\beta, \gamma \in \mathbb{N}^{n-2}$  such that:

$$|\beta| = |\alpha| - 2\alpha_1; \quad |\gamma| = |\alpha| - 2\alpha_2,$$

and

$$(2\alpha_1, 0, \beta) + (0, 2\alpha_2, \gamma) = (2\alpha_1, 2\alpha_2, \beta + \gamma) = 2\alpha.$$

So, for instance, take  $\beta = (\beta_3, \beta_4, \dots, \beta_n)$ ,  $\gamma = (\gamma_3, \gamma_4, \dots, \gamma_n)$ , defined by

$$\beta := (\alpha_3 + \alpha_2 - \alpha_1, \alpha_4, \dots, \alpha_n), \quad \gamma := (\alpha_3 + \alpha_1 - \alpha_2, \alpha_4, \dots, \alpha_n).$$

By construction, we have  $4\alpha_1 + 2|\beta| = 4\alpha_2 + 2|\gamma| = 2|\alpha| \leq 2r$ , so that both  $y_{4\alpha_1, 0, 2\beta}$  and  $y_{0, 4\alpha_2, 2\gamma}$  are diagonal elements of  $M_r(\mathbf{y})$  with at least one entry equal to 0. Hence, by the induction hypothesis,

$$y_{4\alpha_1, 0, 2\beta} \leq \tau_r, \quad y_{0, 4\alpha_2, 2\gamma} \leq \tau_r.$$

Next, consider the two rows and columns indexed by  $(2\alpha_1, 0, \beta)$  and  $(0, 2\alpha_2, \gamma)$ . The constraint  $M_r(\mathbf{y}) \succeq 0$  clearly implies

$$\tau_r^2 \geq y_{4\alpha_1, 0, 2\beta} \times y_{0, 4\alpha_2, 2\gamma} \geq (y_{2\alpha_1, 2\alpha_2, \beta+\gamma})^2 = y_{2\alpha}^2.$$

Hence,  $y_{2\alpha} \leq \tau_r$ , the desired result.  $\square$

## REFERENCES

1. R. ASH, *Real Analysis and Probability*, Academic Press, San Diego, 1972.
2. C. BERG, The multidimensional moment problem and semi-groups, *Proc. Symp. Appl. Math.* **37** (1980), 110–124.
3. G. BLEKHERMAN, There are significantly more nonnegative polynomials than sums of squares, Department of Mathematics, University of Michigan, Ann Arbor, USA, 2004.
4. T. JACOBI, A. PRESTEL, Distinguished representations of strictly positive polynomials, *J. Reine. Angew. Math.* **532** (2001), 223–235.
5. J.B. LASSERRE. Global optimization with polynomials and the problem of moments, *SIAM J. Optim.* **11** (2001), 796–817.
6. J.B. LASSERRE. Polynomials nonnegative on a grid and discrete optimization, *Trans. Amer. Math. Soc.* **354** (2002), 631–649.
7. J.B. LASSERRE. Semidefinite programming vs. LP relaxations for polynomial programming, *Math. Oper. Res.* **27** (2002), 347–360.
8. A.E. NUSSBAUM, Quasi-analytic vectors, *Arch. Mat.* **6** (1966), 179–191.
9. P.A. PARRILO. Structured Semidefinite Programs and Semialgebraic Geometry Methods in Robustness and Optimization, PhD thesis, California Institute of Technology, Pasadena, CA, 2000.
10. P.A. PARRILO. Semidefinite programming relaxations for semialgebraic problems, *Math. Progr. Ser. B* **96** (2003), 293–320.
11. A. PRESTEL, C.N. DELZELL, *Positive Polynomials*, Springer, Berlin, 2001.
12. M. PUTINAR. Positive polynomials on compact semi-algebraic sets, *Indiana Univ. Math. J.* **42** (1993), 969–984.
13. R.T. ROCKAFELLAR, *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.
14. C. SCHEIDERER, Positivity and sums of squares: A guide to some recent results, Department of Mathematics, University of Duisburg, Germany.
15. K. SCHMÜDGEN, The  $K$ -moment problem for compact semi-algebraic sets, *Math. Ann.* **289** (1991), 203–206.
16. M. SCHWEIGHOFER, Optimization of polynomials on compact semialgebraic sets, *SIAM J. Optim.*, to appear.
17. L. VANDENBERGHE AND S. BOYD, Semidefinite programming, *SIAM Review* **38** (1996), pp. 49–95.